Tobacco Use and Help with Quitting Among Adolescents

eyefinity | EHR

In this chapter:

- Denominator, 1
- Numerator, 1
- Eyefinity EHR Instructions, 2

The percentage of adolescents 12 to 20 years of age with a primary care visit during the measurement year for whom tobacco use status was documented and received help with quitting if identified as a tobacco user.

Quality	402
NQF	—
CMS	—

This measure is to be reported once per performance period for patients seen during the performance period. This measure is intended to reflect the quality of services provided for preventive screening for tobacco use.

Denominator

Patients aged 12–20 years;

AND who had one or more visits during the measurement period:

90791, 90792, 90832, 90834, 90837, 90839, 90845, 92002, 92004, 92012, 92014, 96150, 96151, 96152, 97165, 97166, 97167, 97168, 99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99406, 99407, G0438, G0439

Numerator

Patients who were screened for tobacco use at least once within 18 months (during the measurement period or the six months prior to the measurement period);

AND who received tobacco cessation intervention if identified as a tobacco user.

Performance Met

Patient documented as tobacco user AND received tobacco cessation intervention (must include at least one of the following: advice given to quit smoking or tobacco use, counseling on the benefits of quitting smoking or tobacco use, assistance with or referral to external smoking or tobacco cessation support programs, or current enrollment in smoking or tobacco use cessation program) if identified as a tobacco user:

G9458

OR

Current tobacco non-user:

G9459

Performance Not Met

Tobacco assessment OR tobacco cessation intervention not performed, reason not given:

G9460

Eyefinity EHR Instructions

Screen the patient for tobacco use:

1. Open the **Clipboard**.
2. Tap **Edit**.
3. Tap **Social History**.
4. Select a **Smoking Status**.

   If the status is former smoker or nonsmoker, the measure is met; otherwise, compete the following steps.
5. Open the **Ocular Exam**.
6. Select a tobacco use disorder diagnosis from **Impressions**.
7. Select a counseling **Plan**.

8. Select the **Smoking Cessation Counseling** section check box.



9. Tap **Done**.
10. Document the rest of the exam and **Finalize**.

Alternatively, code quality with a special plan:

1. Open the **Ocular Exam**.
2. Select a **Diagnosis** and **Plan**.
3. Tap **Special Plans** and select **MIPS**.
4. From Popular Plans, select **MIPS Quality**.
5. Tap the **More** tab.
6. Select **TOB**.
7. Select the appropriate procedure from the **Quality 402** drop-down.
8. Tap **Done**.
9. Document the rest of the exam and **Finalize**.